

Los Olivos Women's Medical Group

Vaginal Birth After Cesarean (VBAC) Consent

As you know, there are risks to all pregnancies and deliveries. There are specific risks to vaginal delivery after a previous cesarean section (VBAC). Patients should NOT attempt VBAC if they have a previous classical cesarean section, multiple gestation, breech, OR IF THE PATIENT IS UNWILLING TO ASSUME THE ADDED RISKS ASSOCIATED WITH THE TRIAL OF LABOR FOR THEMSELVES AND THEIR BABY. We want to be certain that you clearly understand these risks. Please initial each statement below after you read it to indicate your understanding.

Witness	MD:
Patient	gnature: Date:
	ave read and understand all of the above information. My questions have been answered d I have received all the information I need to make an informed choice after discussing my tions with my doctor.
	nderstand that my doctor may not be available when I go into labor. I understand that some ctors do not perform VBAC deliveries. If my doctor is not immediately available when I go o labor, other alternatives may be recommended including immediate cesarean section by e doctor on call or transfer of care to an Obstetric hospital based physician.
	e risks to me of uterine rupture include but are not limited to hysterectomy, blood nsfusion, infection, injury to bowels, bladder, ureter, blood coagulation defects or death.
	e frequency of death or permanent brain damage after uterine rupture has been reported to as high as 50%, so there is up to a 1 in 200 chance of permanent brain damage or death of baby during a VBAC.
	nderstand that VBAC is associated with a higher risk to my baby than to me. If my uterus otures during VBAC, I understand that no matter how rapidly a cesarean section is done, my by may have permanent brain damage or may die.
	nderstand that the risk of uterine rupture during a VBAC is at least 1%. I understand that if I bose a VBAC and end up having a cesarean during labor, I have a greater risk of problems in if I have an elective repeat cesarean.
	nderstand that approximately 60-70% of women who attempt VBAC have a vaginal delivery. Inderstand that VBAC carries a lower risk to me than a cesarean delivery. The benefits include creased blood loss, decreased post delivery complications and a shorter recuperation.

University HealthCare Alliance ("UHA") is a medical foundation affiliated with Stanford Health Care and Stanford Medicine. UHA contracts with a number of physician groups to provide the medical care in the UHA clinics. Neither UHA, Stanford Health Care, nor Stanford University employ the physicians in the clinics and do not exercise control over the professional services provided by the physician groups.