

## COLORECTAL CANCER SCREENING

### **What is colorectal cancer?**

Colorectal (large bowel) cancer is a disease in which malignant (cancer) cells form in the inner lining of the colon or rectum. Together, the colon and rectum make up the large bowel or large intestine. The large intestine is the last segment of the digestive system (the esophagus, stomach, and small intestine are the first three sections). The large bowel's main job is to reabsorb water from the contents of the intestine so that solid waste can be expelled into the toilet. The first several feet of the large intestine is the colon and the last 6 inches is the rectum.

Most colon and rectal cancers originate from benign wart-like growths on the inner lining of the colon or rectum called polyps. Not all polyps have the potential to transform into cancer. Those that do have the potential are called adenomas. It takes more than 10 years in most cases for an adenoma to develop into cancer. This is why some colon cancer prevention tests are effective even if done at 10-year intervals. This 10-year interval is too long, in some cases, such as in persons with ulcerative colitis or Crohn's colitis, and in persons with a strong family history of colorectal cancer or adenomas.

### **How common is colorectal cancer?**

Colorectal cancer is the second most common cancer killer overall and third most common cause of cancer-related death in the United States in both males and females. Lung and prostate cancers are more common in men and lung and breast in women. In 2010, there were 143,000 new cases and 52,000 deaths from colorectal cancer.

### **What is screening for colorectal cancer?**

Screening means looking for cancer or polyps when patients have no symptoms. Finding colorectal cancer before symptoms develop dramatically improves the chance of survival. Identifying and removing polyps before they become cancerous actually prevents the development of colorectal cancer.

### **Who is at risk for colorectal cancer?**

The average age to develop colorectal cancer is 70 years, and 93% of cases occur in persons 50 years of age or older. Current recommendations are to begin screening at age 50 if there are no risk factors other than age for colorectal cancers. A person whose only risk factor is their age is said to be at average risk. Men tend to get colorectal cancer at an earlier age than women, but women live longer so they 'catch up' with men and thus the total number of cases in men and women is equal. If a person has a history of two or more first-degree relatives (parent, sibling, or child) with colorectal cancer, or any first-degree relatives diagnosed under age 60, the overall colorectal cancer risk is three to six times higher than that of the general population. For those with one first-degree relative diagnosed with colorectal cancer at age 60 or older, there is an approximate two times greater risk of colon cancer than that observed in the general population. Special screening programs are used for those with a family history of colorectal cancer. A well-documented family history of adenomas is also an important risk factor. Persons who have had colorectal cancer or adenomas removed are at increased risk of developing additional adenomas or cancers. Women diagnosed with uterine or ovarian cancer before age 50 are at increased risk of colorectal cancer. These groups should be checked by

colonoscopy at regular intervals, usually every 3 to 5 years. Woman with a personal history of breast cancer have only a very slight increase in risk of colorectal cancer.

### **What are the symptoms of colorectal cancer?**

Symptoms of colorectal cancer vary depending on the location of the cancer within the colon or rectum, though there may be no symptoms at all. The prognosis tends to be worse in symptomatic as compared to asymptomatic individuals. The most common presenting symptom of colorectal cancer is rectal bleeding. Cancers arising from the left side of the colon generally cause bleeding, or in their late stages may cause constipation, abdominal pain, and obstructive symptoms. On the other hand, right-sided colon lesions may produce vague abdominal aching, but are unlikely to present with obstruction or altered bowel habit. Other symptoms such as weakness, weight loss, or anemia resulting from chronic blood loss may accompany cancer of the right side of the colon. You should promptly see your doctor when you experience any of these symptoms.

### **Why should you get checked for colorectal cancer even if you have no symptoms?**

Adenomas can grow for years and transform into cancer without producing any symptoms. By the time symptoms develop; it is often too late to cure the cancer, because it may have spread. Screening identifies cancers earlier and actually results in cancer prevention when it leads to removal of adenomas (pre-cancerous polyps).

### **What else can I do to prevent the development of colorectal cancer?**

The strategy for reducing colorectal cancer deaths is simple.

For normal risk individuals, screening tests begin at age 50 and the preferred approach is a screening colonoscopy every 10 years; an alternate strategy consists of annual stool test for blood and a flexible sigmoidoscopic exam every 3 to 5 years. Colonoscopic surveillance (also called screening colonoscopy) needs to be available at more frequent intervals for individuals at high risk for colon cancer (for instance, those with a personal history of colorectal cancer or adenomatous polyps; family history of colorectal cancer; non-hereditary polyposis; colorectal cancer; or a pre-disposing condition such as inflammatory bowel disease. For both average and high risk individuals, all potential pre-cancerous polyps must be removed.

Recent observations suggest regular use of non-steroidal anti-inflammatory drugs or aspirin reduce the chances of colorectal cancer death by 30-50%. These drugs also have risks, particularly intestinal bleeding, and patients should consult their physician as to whether regular use of these agents is appropriate. Folate, calcium, and post-menopausal estrogens each have a modest protective benefit against colon cancer. A high fiber (vegetables) and low fat diet, regular exercise, maintenance of normal body weight and cessation of smoking are also beneficial. None of the measures is as effective as, or should replace, colorectal cancer screening.

### **Gastroenterologists**

Marwan Balaa, MD	402-9990
Alfred Hurwitz, MD	294-4272
Michael Kushlan, MD	358-0468
Terry Lin, OD	358-6580
Jay Raju, MD	297-2416
Bassam Saffouri, MD	370-7801
Geoffrey Spenser, MD	626-7375
David Stein, MD	626-7375
Kevin Stuart, MD	847-1311

### **Colorectal Surgeons**

Richard Coughlin, MD	358-2868
Kristina Hobson, MD	358-2868
Mark Segall, MD	358-3500
Peter Youn, MD	358-2868